Saskatchewan
Ministry of
Agriculture

Crop Report

Crop Report for the Period August 24 to 30, 2010

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 22, September 2, 2010

Eight per cent the 2010 crop has been combined, according to Saskatchewan Agriculture's weekly Crop Report. An additional 22 per cent is swathed or ready to straight-combine. The five-year provincial average (2005 to 2009) for this time of year is 28 per cent combined and 25 per cent swathed or ready to straight-combine. Another wet week in most areas of the province has delayed crop maturity and slowed harvest.

The southwest has 15 per cent of the crop combined, the southeast 14 per cent, and the northeast and northwest have two per cent combined. In east-central Saskatchewan, four per cent of the crop is combined and, in west-central Saskatchewan, the figure is six per cent.

Haying is wrapping up in most regions. The unsettled weather has extended the haying season for Saskatchewan producers.

Cropland topsoil moisture is rated as 29 per cent surplus, 68 per cent adequate and three per cent short. Topsoil moisture on hay land and pasture is rated as 17 per cent surplus, 78 per cent adequate and five per cent short.

The majority of crop damage is being caused by rain, wind and disease. The strong wind in some areas last week caused some canola, pea and hay swaths to blow around.

Farmers are busy haying, getting ready for harvest, harvesting and hauling bales. Winter wheat is being seeded in some areas.

One year ago
Just under eight per cent of the 2009 crop had been combined. An additional 13 per cent was swathed or ready to straight combine. The week was good maturing and harvesting weather.

Southeastern Saskatchewan (Crop Districts 1, 2, and 3ASE)

Farmers were able to get some combining done during the week, however, by the weekend, the rain set in again, slowing harvest progress. The amount of rain falling on the region ranged from nil to 110 mm. The majority fell on Sunday and Monday (Aug 29 and 30). Windthorst was hit extremely hard, recording 110 mm of rain. The Kisbey area reported 31 mm, the Kennedy and

For further information, contact: Grant McLean, PAg,
Cropping Management Specialist, Agriculture Knowledge Centre,
Toll Free: 1-866-457-2377, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

Tantallon areas 50 mm, the Fillmore and Griffin areas 35 mm, the Moose Jaw area 27 mm and the Ceylon area 36 mm. Most areas reported more than 20 mm of rain.

Farmers made good harvest progress at the beginning of the week and have 14 per cent of the crop combined and 30 per cent of the crop swathed or ready to straight-cut. Last year at this time, eight per cent of the crop had been combined and 17 per cent swathed or ready to straight-cut.

Many crop reporters are having difficulty assessing yields at this time due to crop immaturity, flooded areas and disease. The average yields for the region are as follows: winter wheat 50 bu/ac, spring wheat 34 bu/ac, durum 32 bu/ac, oats 65 bu/ac, barley 51 bu/ac, fall rye 48 bu/ac, flax 20 bu/ac, canola 28 bu/ac, mustard 880 lb./ac, lentils 1,100 lb./ac, peas 30 bu/ac, canaryseed 1,400 lb./ac and chickpeas 1,200 lb./ac.

Haying has essentially finished in the region. Farmers are cutting greenfeed crops.

Topsoil moisture conditions on cropland are reported as 41 per cent surplus, 58 per cent adequate and one per cent short. On hay land and pasture, topsoil moisture is rated as 19 per cent surplus, 78 per cent adequate and three per cent short. Seventy-three per cent of the cropland in Crop District 1B has surplus topsoil moisture.

Crop damage is mostly attributed to wind (lodging and windblown swaths), rain and flooding. Disease is also causing crop damage. The rain is causing some concerns about disease in ripening pulse crops. Sprouting and bleaching in swathed cereals is also a concern.

Although there was one day when the temperature was above 30C, several days of warm weather are still needed to push the crops along. Many areas are still one to two weeks behind normal crop development. Wet fields are causing problems with harvesting equipment. Some farmers are having to dump grain into grain carts on the road as the fields are too soft. With all the rain, many crop reporters are concerned about crop quality. Heavy dew in the mornings and again in the evenings makes for short combining days. Very little fall seeding is being done. Farmers are haying, harvesting, getting ready for harvest and controlling weeds on unseeded acres.

Southwestern Saskatchewan (Crop Districts 3ASW, 3AN, 3B and 4)

The region received between zero and 38 mm of rain during the week. Most of it fell on Sunday and Monday. The Viceroy area received 35 mm, the Fife Lake area 32 mm, the Coderre area 24 mm, the Mossbank area 20 mm, the Shaunavon area 30 mm, the Vanguard area 5 mm, the Rush Lake area 17 mm, the Gull Lake area 38 mm, the Eastend area 25 mm, the Hazlet area 21 mm and the Maple Creek and Fox Valley areas 15 mm. Most of the region received more than 15 mm.

Good harvest progress was made until the rain late in the week. The region has 15 per cent of the 2010 crop combined and an additional 16 per cent swathed or ready to straight-cut. Last year at this time, 26 per cent of the crop had been combined and 17 per cent swathed or ready to straight-cut.

Crop reporters are indicating the following average crop yields at this time: winter wheat 41 bu/ac, spring wheat 32 bu/ac, durum 33 bu/ac, oats 50 bu/ac, barley 48 bu/ac, fall rye 54 bu/ac, triticale 35 bu/ac, flax 20 bu/ac, canola 27 bu/ac, mustard 1,000 lb./ac, lentils 1,200 lb./ac, peas 30 bu/ac, canaryseed 1,300 lb./ac and chickpeas 1,000 lb./ac.

Haying is wrapping up in the region.

Cropland topsoil moisture is rated as 19 per cent surplus, 79 per cent adequate and two short. Hay land and pasture topsoil moisture conditions are rated as six per cent surplus, 91 per cent adequate and three per cent short.

Crop damage has been caused by rain, disease and wind (lodging and windblown swaths). Sawflies are causing damage in the wheat, making some farmers hesitant to swath due to the rain.

Farmers are busy finishing up haying, harvesting, getting ready for harvest and hauling bales. Frost is a worry as temperatures have dipped to 5C on occasion. Rain on desiccated/swathed pulses and cereals has farmers worried about crop quality. A few weeks of dry and sunny weather are needed, as most crops are one to two weeks behind normal in development.

East-Central Saskatchewan (Crop Districts 5 and 6a)

It was another wet week in the east-central region. Rain fell in amounts ranging from zero to 58 mm. CD 5A received between 8 mm and 64 mm. CD 5B received between 2 mm and 28 mm. CD 6A received between zero and 31 mm. Since April 1, the region has received between 271 mm and 752 mm. The Ituna area recorded 58 mm, the Goodeve area 8 mm, the Churchbridge area 36 mm, the Kamsack area 28 mm, the Quill Lake area 2 mm, the Craven area 31 mm and the Leroy area 10 mm.

The weather co-operated early in the week and allowed some harvesting and haying to continue. Four per cent of the crop is combined and an additional 26 per cent is swathed or ready to straight-cut. Last year at this time, two per cent of the crop had been combined and eleven per cent swathed or ready to straight-cut.

Crop yields are difficult to assess at this time. Crop reporters are indicating the following average yields: winter wheat 42 bu/ac, spring wheat 32 bu/ac, durum 34 bu/ac, oats 65 bu/ac, barley 50 bu/ac, fall rye 30 bu/ac, triticale 25 bu/ac, flax 26 bu/ac, canola 24 bu/ac, mustard 900 lb./ac, lentils 1,100 lb./ac, peas 32 bu/ac, canaryseed 900 lb./ac and chickpeas 1,400 lb./ac.

Haying is continuing in the region. Many producers are indicating that quality is poor because of all the rain that has fallen on the swaths. Some hay fields are under water.

Topsoil moisture on cropland is reported as 59 per cent surplus, 40 per cent adequate and one per cent short. Hay land and pasture topsoil moisture is rated as 50 per cent surplus, 47 per cent adequate and three per cent short. In CDs 5A and 5B respectively, 75 and 89 per cent of the cropland has surplus topsoil moisture.

Flooding, wind (lodging) and rain were the major causes of crop damage. Rain on mature crops is leading to damage from crop diseases. The number of flooded acres continues to increase as the rain continues.

Farmers are busy haying, getting ready for harvest, harvesting and controlling weeds on unseeded acres. Canola crops are weedy in some areas. Some crops are ready to swath, but the rain is keeping farmers out of the fields. Crop quality is being reduced in the swath. Some winter wheat is being seeded, although most fields are difficult to access with seeding equipment. There are reports of combines getting stuck in winter wheat fields. In general, crops need two to three weeks to mature. Sunny and dry weather is needed. Frost is a concern.

West-Central Saskatchewan (Crop Districts 6B and 7)

The rain missed the majority of the west-central region, so good harvest progress was made. The amount of rain that fell in the region ranged from zero to 11 mm. The Dinsmore area recorded 11 mm, the Elbow and Conquest areas 3 mm and the Outlook, Hanley and Kerrobert areas 1 mm. One crop reporter indicated this is the first week since April 1 in which he has not recorded rain.

Six per cent of the 2010 has been combined, with an additional 18 per cent swathed or ready to straight-cut. Farmers were focused on swathing and desiccating this week. Last year at this time, four per cent of the crop was combined and 16 per cent was swathed or ready to straight-combine.

Crop reporters are indicating the average crop yields at this time are as follows: winter wheat 43 bu/ac, spring wheat 37 bu/ac, durum 39 bu/ac, oats 76 bu/ac, barley 56 bu/ac, fall rye 54 bu/ac, triticale 42 bu/ac, flax 25 bu/ac, canola 30 bu/ac, mustard 1,200 lb./ac, lentils 1,200 lb./ac, peas 37 bu/ac and canaryseed 1,200 lb./ac.

Haying operations are wrapping up in the region. Good haying progress was made this week.

Cropland topsoil moisture is rated as 15 per cent surplus, 80 per cent adequate and five per cent short. Hay land and pasture topsoil moisture conditions are rated as five per cent surplus, 88 per cent adequate and seven per cent short.

The majority of crop damage is due to disease. The moisture received in the past few weeks has farmers worried about crop quality. Pulses, particularly lentils, are showing high disease pressures. High winds scattered some swaths.

Farmers are busy haying, getting ready for harvest and harvesting. Crops are behind normal in development by one to two weeks, so many farmers are hoping for warm and sunny weather. Farmers are swathing peas, canola and winter cereals, and desiccating lentils and peas.

Northeastern Saskatchewan (Crop Districts 8 and 9AE)

Most of the region received no rain for the week, although the Porcupine Plain area reported 23 mm, the Tisdale area 7 mm and the Alvena area 2 mm.

Significant harvest progress was made this week. Two per cent of the 2010 crop is combined and an additional 25 per cent is swathed or ready to straight-cut.

Crop yield predictions are quite variable across all crops. Crop reporters in the region are indicating the following average crop yields at this time: winter wheat 38 bu/ac, spring wheat 33 bu/ac, oats 68 bu/ac, barley 50 bu/ac, fall rye 37 bu/ac, flax 17 bu/ac, canola 24 bu/ac, peas 27 bu/ac and canaryseed 1,000 lb./ac.

Good haying progress was made this past week and most producers are nearly finished.

Cropland topsoil moisture is rated as 37 per cent surplus and 63 per cent adequate. Hay land and pasture topsoil moisture is rated as 26 per cent surplus, 67 per cent adequate and seven per cent short. Seventy-two per cent of the cropland in CD 8A has surplus topsoil moisture.

Crop damage is due to disease and wind (lodging).

Farmers are busy haying, getting ready for harvest, harvesting and controlling weeds on unseeded acres. Some unseeded fields have been sprayed four times this year. Crop staging is variable, and crops are still one to two weeks behind normal in development. Fields are very wet in areas and swathers are getting stuck.

Northwestern Saskatchewan (Crop Districts 9AW and 9B)

Only a few areas in the region received any moisture this week. The Rapid View area received 19 mm, the Meadow Lake and Dorintosh areas 2 mm and the Barthel area 3 mm. Low temperatures were recorded in some areas. The Neilburg area hit 0C and the Glaslyn area 3C.

Significant progress was made with swathing. Two per cent of the crop is combined and an additional 18 per cent is swathed or ready to straight-cut.

Estimated average crop yields for the region are as follows: spring wheat 38 bu/ac, oats 79 bu/ac, barley 65 bu/ac, fall rye 32 bu/ac, flax 20 bu/ac, canola 32 bu/ac and peas 38 bu/ac.

Good haying progress was made, and most producers are nearly finished although there are some areas in which farmers still have some mowing and baling to do.

Cropland topsoil moisture is rated as one per cent surplus, 93 per cent adequate and six per cent short. Hay land and pasture topsoil moisture is rated as 92 per cent adequate and eight per cent short.

Wind and waterfowl are causing crop damage. The wind blew crop and hay swaths around. Disease pressures in pulses have caused some crop damage as well. Some crops are lodging in areas that received heavy rain over the past couple of weeks.

Warm and sunny weather is needed. Farmers are busy finishing haying, hauling bales, getting ready for harvest and harvesting.

Winter wheat	% Standing	% in swath	% ready to straight combine	% combined
SE	7	4	11	78
SW*	14	3	15	68
EC	15	6	34	45
WC	50	20	8	22
NE	45	15	5	35
NW	90	10	0	0

Fall rye	% Standing	% in swath	% ready to straight combine	% combined
SE	10	1	6	83
SW*	12	12	5	60
EC	9	15	19	57
WC	3	63	5	29
NE	20	15	25	40
NW	82	18	0	0
* 11 per cent is rated as 'other'				

Spring wheat	% Standing	% in swath	% ready to straight combine	% combined
SE	69	8	17	6
SW*	90	2	6	2
EC	93	2	4	1
WC	93	4	3	0
NE	95	3	1	1
NW	95	4	1	0

Durum	% Standing	% in swath	% ready to straight combine	% combined
SE	83	6	9	2
SW	90	2	6	2
EC	98	2	0	0
WC	93	6	1	0
NE	90	9	0	1
NW	N/A	N/A	N/A	N/A

Oats	% Standing	% in swath	% ready to straight combine	% combined
SE	86	8	2	4
SW*	82	7	2	2
EC**	98	1	0	0
WC	96	4	0	0
NE	95	5	0	0
NW	91	6	0	0
*seven per cent is rated as 'other** one per cent is rated as 'other" three per cent is rated as 'other'				

Barley	% Standing	% in swath	% ready to straight combine	% combined
SE	65	12	8	15
SW	91	1	5	3
EC	89	5	4	2
WC	96	4	0	0
NE	81	8	7	4
NW	86	13	1	0

Lentils	% Standing	% in swath	% ready to straight combine	% combined
SE	40	18	9	33
SW	46	7	23	24
EC	55	2	37	6
WC	84	9	6	1
NE	63	10	26	1
NW	81	0	3	16

Peas	% Standing	% in swath	% ready to straight combine	% combined
SE	16	12	11	61
SW	21	0	16	63
EC	36	6	31	27
WC	66	6	20	8
NE	61	4	22	13
NW	72	2	11	15

Canola	% Standing	% in swath	% ready to straight combine	% combined
SE	33	57	0	10
SW	32	56	6	6
EC	53	45	1	1
WC	56	44	0	0
NE	55	44	0	1
NW	65	35	0	0

Mustard	% Standing	% in swath	% ready to straight combine	% combined
SE	66	27	1	6
SW	53	12	18	17
EC	66	18	9	7
WC	77	22	0	1
NE	95	5	0	0
NW	100	0	0	0

Triticale	% Standing	% in swath	% ready to straight combine	% combined
SE	100	0	0	0
SW*	48	9	2	4
EC	100	0	0	0
WC	96	4	0	0
*37 per cent is rated as 'other'.				

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

for the week ending August 30, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010

Weekly Rainfall

for the week ending August 30, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010

Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period to August 24 to 30, 2010

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	1	Argyle	17.0	436.0
	2	Mount Pleasant	9	328
	3	Enniskillen	9	352
	33 A	Moose Creek	N/A	249
	33 B	Moose Creek	15	385
	34	Browning	27	347
	61	Antler	N/A	490
	63	Moose Mountain	13	410
	64	Brock	31	393
	65	Tecumseh	26	401
1B	91	Maryfield	16	411
	92	Walpole	7	358
	93	Wawken	50	480
	95	Golden West	110	452
	122	Martin	12	412
	123 A	Silverwood	18	326
	123 B	Silverwood	N/A	N/A
	124	Kingsley	25	345
	125	Chester	14.6	356
	151 A	Rocanville	50	491
	151 B	Rocanville	N/A	480
	153	Willowdale	38	487
	154	Elcapo	N/A	425
	155	Wolseley	35	626
2A	66	Griffin	35	514
	67	Weyburn	22	456
	68	Brokenshell	25	386
	69	Norton	N/A	313
	96	Fillmore	35	355
	97	Wellington	22	332
2B	127 A	Francis	12	391
	127 B	Francis	NIL	268
	129	Bratt's Lake	3	336
	156 A	Indian Head	9	459
	156 B	Indian Head	12	450
	157	South Qu'appelle	N/A	NIL
	160	Pense	26	370
	161	Moose Jaw	27	379
	191	Marquis	23	429
3ASE	38 A	Laurier	N/A	423
	38 B	Laurier	36	398
	39 A	The Gap	24	345
	39 B	The Gap	23	490
3ASW	10	Happy Valley	25	410
	11	Hart Butte	N/A	NIL
	12	Poplar Valley	32	299
	40	Bengough	29	382
	42	Willow Bunch	21	352
	43	Old Post	21	351
	44	Waverley	9	369
	70	Key West	25	371
	71 A	Excel	35	378
	71 B	Excel	16	288
	73 A	Stonehenge	NIL	315
	73 B	Stonehenge	23	423
	74	Wood River	26	348
3AN	101	Terrell	17	405
	102	Lake Johnston	20	322
	103	Sutton	20	344
	134	Shamrock	24	434
	164	Chaplin	18	375
	193	Eyebrow	12	382
	194	Enfield	15	411
3BS	17	Val Marie	17	20
	45	Mankota	20	292
	75 A	Pinto Creek	2	308
	75 B	Pinto Creek	17	308
	76 A	Auvergne	9	276
	76 B	Auvergne	13	266
	77	Wise Creek	8	275
	78 A	Grassy Creek	30	390
	78 B	Grassy Creek	23	370
	78 C	Grassy Creek	N/A	361
	105	Glenbain	19	303
	106	Whiska Creek	5	241
	107	Lac Pelletier	12	308
	108	Bone Creek	34	395
3BN	138	Webb	N/A	372
	165	Morse	N/A	NIL
	166	Excelsior	17	489
	167	Sask. Landing	11	437
	168	Riverside	14	450
	226	Victory	N/A	402
	228	Lacadena	14	449
	255	Coteau	N/A	NIL
	257	Monet	8	400

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	49	White Valley	20	353
	49 A	White Valley	25	294
	51	Rano	15	373
	79	Arlington	27	410
	109	Carmichael	38	563
	110	Piapot	18	421
	111	Maple Creek	25	420
4B	139	Gull Lake	18	517
	141	Big Stick	15	443
	142	Enterprise	7	523
	169	Pittville	21	403
	171	Fox Valley	15	399
	229	Miry Creek	18	515
	231	Happyland	5	415
5A	181	Langenberg	36	433
	183	Fertile Belt	35	486
	185	Mcleod	27	435
	186	Abernethy	15	357
	211	Churchbridge	21	442
	213	Saltcoats	33	507
	216	Tullymet	58	288
	241	Calder	38	434
	243	Wallace	31	404
	245 A	Garry	12	494
	245 B	Garry	8	520
	246	Ituna Bon Accord	10	407
	247	Kelross	21	511
	248	Touchwood	11	277
5B	271	Cote	28	435
	275	Insinger	N/A	NIL
	276	Foam Lake	7	645
	277	Emerald	11	617
	278	Kutawa	N/A	NIL
	305	Invermay	8	752
	307	Elfros	26	748
	308	Big Quill	6	480
	333	Clayton	N/A	134
	335	Hazel Dell	10	468
	336	Sasman	21	629
	337	Lakeview	3	490
	338	Lakeside	2	514
	366	Kelvington	8	423
6A	190 A	Dufferin	8	271
	190 B	Dufferin	24	388
	190 C	Dufferin	6	322
	190 D	Dufferin	N/A	339
	219 A	Longlaketon	25	360
	219 B	Longlaketon	31	418
	220	Mckillop	5	344
	221	Samia	20.8	436
	222	Craik	13	454
	251	Big Arm	6	430
	252	Arm River	12	367
	279	Mount Hope	22	449
	280	Wreford	2	387
	282	McCraney	2	436
	309	Prairie Rose	14	481
	313	Lost River	NIL	401
	339	Leroy	10.4	529
	340	Wolverine	NIL	516
	341	Viscount	N/A	489
	343	Blucher	NIL	509
6B	253	Willner	3.5	334
	283	Rosedale	NIL	505
	284	Rudy	1	482
	285	Fertile Valley	3	443
	286	Milden	11	468
	314	Dundurn	1	509
	344	Corman Park	NIL	464
	346	Perdue	NIL	524
	376 A	Eagle Creek	NIL	633
	376 B	Eagle Creek	NIL	463
7A	260	Newcombe	NIL	390
	260 B	Newcombe	N/A	293
	287	St. Andrews	NIL	365
	288	Pleasant Valley	N/A	383
	290 A	Kindersley	N/A	385
	290 B	Kindersley	N/A	390
	292	Milton	N/A	334
	317 A	Marriott	NIL	522
	317 B	Marriott	NIL	548
	318 A	Mountain View	NIL	501
	318 B	Mountain View	NIL	515
	320	Oakdale	NIL	363
	321	Prairiedale	NIL	386

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7B	347 A	Biggar	NIL	408
	347 B	Biggar	N/A	415
	350 A	Mariposa	NIL	305
	350 B	Mariposa	1	381
	351	Progress	N/A	414
	352	Heart's Hill	NIL	460
	377	Glenside	N/A	463
	378 A	Rosemount	NIL	348
	378 B	Rosemount	NIL	439
	379	Reford	NIL	397
	380	Tramping Lake	N/A	350
	381	Grass Lake	N/A	506
	382	Eye Hill	NIL	376
	409	Buffalo	N/A	525
	410	Round Valley	NIL	437
	411	Senlac	N/A	431
8A	395	Porcupine	23	461
	397	Barrier Valley	N/A	496
	428	Star City	N/A	429
	456	Arborfield	4	400
	457	Connaught	7	576
	458	Willow Creek	5	462
	486	Moose Range	NIL	412
8B	369	St. Peter	NIL	481
	370	Humboldt	NIL	575
	371	Bayne	NIL	515
	372	Grant	NIL	496
	402	Fish Creek	2	424
	429 A	Flett's Springs	NIL	319
	429 B	Flett's Springs	NIL	459
	430	Invergordon	N/A	43
	459	Kinistino	NIL	425
9AE	481	Prince Albert	NIL	442
	488	Torch River	NIL	453
	520	Paddockwood	N/A	67
9AW	406	Mayfield	NIL	334
	435	Redberry	NIL	454
	436	Douglas	NIL	252
	463	Duck Lake	NIL	406
	464	Leask	NIL	452
	467	Round Hill	N/A	NIL
	494	Canwood	N/A	446
	496	Spiritwood	NIL	526
	497	Medstead	N/A	374
9B	438	Battle River	NIL	407
	440	Hillsdale	NIL	360
	442	Manitou Lake	NIL	335
	498 A	Parkdale	NIL	370
	498 B	Parkdale	NIL	319
	499	Mervin	NIL	315
	501 A	Frenchman Butte	NIL	354
	501 B	Frenchman Butte	NIL	309
	501 C	Frenchman Butte	N/A	433
	502	Britannia	NIL	442
	561	Loon Lake	3	352
	588 A	Meadow Lake	2	379
	588 B	Meadow Lake	19	403
	588 C	Meadow Lake	3	410
	588 D	Meadow Lake	2	384
	622	Beaver River	NIL	356

Municipality No: A, B, C, and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2010
To: August 30, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200
Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010

Cumulative Rainfall

From: April 1, 2010
To: August 30, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010

Cropland Topsoil Moisture Conditions

August 31, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010

Hay and Pasture Topsoil Moisture Conditions

August 31, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010

Cropland Topsoil Moisture Conditions

August 31, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010

Hay and Pasture Topsoil Moisture Conditions

August 31, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: September 1, 2010